Stapled Hemorrhoidopexy for the Treatment of Hemorrhoidal Disease: A Video Vignette

🖲 Ramazan Kozan¹, 🖻 Can Sahin², 🖻 Safa Özaydın¹, 🖻 Sezai Leventoqlu¹

¹Gazi University Faculty of Medicine, Department of Surgery, Ankara, Turkey ²Yıldırım Beyazıt University, Yenimahalle Training and Research Hospital, Clinic of Surgery, Ankara, Turkey

ABSTRACT

Hemorrhoidal disease is a common proctologic disease characterized by enlarged, inflamed, thrombosed, or prolapsed hemorrhoids, with symptoms including pain and rectal bleeding. This video presents a 28-year-old female patient with a grade 3-4 hemorrhoidal disease who underwent stapled hemorrhoidopexy using the prolapse and hemorrhoids system.

Keywords: Hemorrhoids, proctology, surgery

Introduction

Stapled hemorrhoidopexy is an efficient surgical procedure for hemorrhoidal diseases, particularly with mucosal prolapse.1 This technique reduces the length of hospital stays and may have an advantage in terms of decreased operating times, reduced post-operative pain, and less bleeding; however, it is associated with an increased rate of recurrent prolapses.²⁻⁴ Almost all the recurrence cases are related to technical failures. This video aims to show the detailed technique steps of the stapled hemorrhoidopexy to prevent future complications during this procedure.

The procedure for prolapse and hemorrhoids was introduced in 1993 as a novel treatment for hemorrhoidal disease and was originally described as rectal mucosectomy. The procedure's creator, Antonio Longo, described this surgery as an excision of a rectal internal mucosal prolapse.5

Case Report

This video introduces a case of a 28-year-old woman who had hemorrhoids. The patient had complaints of anal swelling, soiling, and bleeding. The physical examination revealed grade 3-4 hemorrhoidal disease. Her obstructed defecation score was 5, and a grade 1 rectocele was identified during a pelvic floor examination. No other issues were found during the rectosigmoidoscopy. A stapled hemorrhoidopexy procedure was performed, and the patient was discharged on postoperative day one and received recommendations for daily wound care. The postoperative visual analog scale scores were 1, 0, 0, and 0 on the first day, first week, first month, and third month, respectively. No recurrence was observed during the 3-month postoperative follow-up.

Ethics

Authorship Contributions

Surgical and Medical Practices: R.K., S.L., Concept: R.K., S.Ö., S.L., Design: R.K., Data Collection or Processing: S.Ö., Analysis or Interpretation: C.Ş., Literature Search: C.Ş., Writing: C.Ş. Conflict of Interest: No conflict of interest was declared by the authors.



Video 1. 10.4274/tjcd.galenos.2023.2023-5-3.video1



Address for Correspondence: Ramazan Kozan, MD, Gazi University Faculty of Medicine, Department of Surgery, Ankara, Turkey E-mail: dr.kozan@hotmail.com ORCID ID: orcid.org/0000-0002-3835-8759

@ (€) (⊜) |

Copyright® 2024 The Author. Published by Galenos Publishing House on behalf of Turkish Society of Colon and Rectal Surgery . This is an open access article under the Creative Commons AttributionNonCommercial 4.0 International (CC BY-NC 4.0) License

Financial Disclosure: The authors declared that this study received no financial support.

References

- Pata F, Sgró A, Ferrara F, Vigorita V, Gallo G, Pellino G. Anatomy, Physiology and Pathophysiology of Haemorrhoids. Rev Recent Clin Trials 2021;16:75-80.
- Puia IC, Puia A, Florea ML, Cristea PG, Stanca M, Fetti A, Moiş E. Stapled Hemorrhoidopexy: Technique and Long Term Results. Chirurgia (Bucur) 2021;116:102-108.
- 3. Nisar PJ, Acheson AG, Neal KR, Scholefield JH. Stapled hemorrhoidopexy compared with conventional hemorrhoidectomy: systematic review of randomized, controlled trials. Dis Colon Rectum 2004;47:1837-1845.
- Tjandra JJ, Chan MK. Systematic review on the procedure for prolapse and hemorrhoids (stapled hemorrhoidopexy). Dis Colon Rectum 2007;50:878-892.
- Eberspacher C, Magliocca FM, Pontone S, Mascagni P, Fralleone L, Gallo G, Mascagni D. Stapled Hemorrhoidopexy: "Mucosectomy or Not Only Mucosectomy, This Is the Problem". Front Surg 2021;8:655257.